

Predictors of inpatient mortality among children hospitalized for severe acute malnutrition: a systematic review and meta-analysis

Radhini Karunaratne,¹ Jonathan P Sturgeon,^{1,2} Rajvi Patel,³ and Andrew J Prendergast^{1,2}

¹Zvitambo Institute for Maternal and Child Health Research, Harare, Zimbabwe; ²Centre for Genomics and Child Health, Blizard Institute, Queen Mary University of London, London, United Kingdom; and ³Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, United Kingdom

ABSTRACT

Background: Malnutrition underlies 45% of under-5 deaths globally. Severe acute malnutrition (SAM) is the most serious form of undernutrition, characterized by wasting with or without edema. Mortality remains high (10%–40%) among children requiring hospitalization for complicated SAM.

Objectives: We aimed to systematically document the factors independently associated with inpatient mortality in children with SAM.

Methods: Embase, Ovid MEDLINE, the Cochrane Library, and clinicaltrials.gov were searched for articles published between January 2000 and January 2020, using a prespecified protocol. Eligible studies included children aged ≤ 59 mo hospitalized with SAM and used multivariable analysis to assess the baseline factors independently associated with inpatient mortality. Random-effects meta-analysis, stratified by the stated measure of effect, was used where $>20\%$ of studies included the same factor in analyses.

Results: Twenty-eight of 1432 studies fulfilled inclusion criteria: 19 studies included all children with SAM and 9 included specific subgroups of children with SAM. All 19 main studies were from 8 countries across Africa, with a median of 400 children/study. The mean inpatient mortality was 15.7% (95% CI: 10.4%, 21.0%) and HIV prevalence ranged from 2.1% to 51%. Nine factors were included in the meta-analysis, stratified by HR and OR. HIV infection (HR: 4.32; 95% CI: 2.31, 8.08), weight-for-height z score (WHZ) (OR: 0.44; 95% CI: 0.24, 0.80), diarrhea (HR: 2.84; 95% CI: 1.40, 5.75), pneumonia (HR: 1.89; 95% CI: 1.19, 3.02), presence of shock (HR: 3.67; 95% CI: 2.24, 6.03), and lack of appetite (HR: 2.16; 95% CI: 1.48, 3.16) were associated with increased mortality, whereas child age and sex were not. The association between edema and mortality was difficult to ascertain from the available studies.

Conclusions: HIV infection, diarrhea, pneumonia, shock, lack of appetite, and lower WHZ are independent predictors of inpatient mortality in children with SAM. These factors may help to risk-stratify children being hospitalized with complicated SAM. This systematic review/meta-analysis protocol was registered at www.crd.york.ac.uk/prospero as CRD42019152267. *Am J Clin Nutr* 2020;00:1–11.

Keywords: malnutrition, SAM, mortality, child, inpatient, severe acute malnutrition, predictors

Introduction

Globally, malnutrition underlies 45% of deaths in children under-5, leading to an estimated 3.1 million deaths per year (1). Severe acute malnutrition (SAM), the most serious manifestation of undernutrition, is a complex syndrome characterized by wasting of tissues with or without edema, together with co-infections, enteropathy, metabolic dysregulation, and inflammation (2, 3). The WHO first published guidance on the inpatient management of “protein energy malnutrition” in 1981 (4). With the advent of 2 formula diets, F-75 and F-100, guidelines were updated in 1999 (5) to define 10 steps of standardized inpatient treatment. The guidelines were further revised in 2003 (6), 2013 (7), and 2016 (8), to reflect the development of ready-to-use therapeutic foods, the availability of the 2006 WHO Child Growth Standards, and the increasing use of midupper arm circumference (MUAC) for diagnosis, which together increased the feasibility of community management.

No specific funding for this study, but the authors are supported by Wellcome Trust grants 108065/Z/15/Z (to AJP) and 220566/Z/20/Z (to JPS).

Supplemental Table 1 and Supplemental Figure 1 are available from the “Supplementary data” link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/ajcn/>.

Data described in the article will be made available upon request pending application and approval.

RK and JPS contributed equally to this work.

Address correspondence to JPS (e-mail: jpsturgeon@cantab.net).

Abbreviations used: aHR, adjusted hazard ratio; aOR, adjusted odds ratio; ART, antiretroviral therapy; CRP, C-reactive protein; MUAC, midupper arm circumference; NG, nasogastric; SAM, severe acute malnutrition; WHZ, weight-for-height z score.

Received April 27, 2020. Accepted for publication June 12, 2020.

First published online 0, 2020; doi: <https://doi.org/10.1093/ajcn/nqaa182>.

Current guidelines classify uncomplicated and complicated SAM according to the absence or presence of medical complications. Currently, only children with complications such as edema, lack of appetite, or infections are hospitalized for nutritional rehabilitation (5). Despite this stratified approach and standardization of care, hospital mortality remains at 10%–40% (9), for reasons that remain unclear. The primary aim of this systematic review and meta-analysis was to identify the baseline factors independently associated with inpatient mortality in children hospitalized with SAM.

Methods

Search strategy and selection criteria

We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines throughout this review. We searched Ovid MEDLINE®, Embase®, the Cochrane Library [the Cochrane Database of Systematic Reviews and the Cochrane Central Register of Controlled Trials (CENTRAL)], and clinicaltrials.gov for articles published in English. Because we were interested in studies published since adoption of the WHO guidelines, articles were included if they were published from 1 January, 2000 until the date of the initial search (5 July, 2019). The searches were repeated on 9 January, 2020 to identify new articles. Full details of the search strategies can be found in the systematic review protocol, which was registered with PROSPERO (CRD42019152267).

Participants were children aged ≤ 59 mo, hospitalized with SAM. Owing to varying classifications over time, SAM was defined using the diagnostic criteria employed by each study. Studies from any setting were included, regardless of the SAM treatment guidelines used. We included all study types, provided they evaluated baseline factors associated with inpatient mortality and used multivariable analysis to control for confounding. We did not prespecify the baseline factors of interest, which could therefore include any physical, biological, demographic, or social factors measured by each study. The primary outcome of interest was inpatient mortality. Studies were excluded if inpatient mortality was not specifically reported. Nonhuman studies were excluded. Studies that specifically investigated a subpopulation of children with SAM were included, but were analyzed separately from studies including all children with SAM.

Data selection and extraction strategy

Initial titles from Ovid MEDLINE® and Embase® were screened by 1 reviewer. Studies that were potentially relevant were deduplicated using Endnote X9 (Clarivate Analytics, 2013), and abstracts were screened by 2 separate reviewers; full-text articles were retrieved for studies likely to meet the inclusion criteria. Articles were assessed by 1 reviewer against the inclusion/exclusion criteria; a random 10% sample were screened independently by a second reviewer. Any discrepancies were resolved by discussion with a third reviewer. The data were extracted into a prepiloted spreadsheet. The bibliographies of all included studies, and of any review articles or protocols found during the search, were examined for additional studies meeting

the inclusion criteria. Data on individual factors associated with inpatient mortality were extracted as presented in each article after multivariable analysis (i.e., adjusted HR, RR, or OR). Where full results were not provided in the article, study authors were contacted to request data.

Assessment of bias

Studies were assessed using the methodology checklist employed for the evaluation of the quality of prognostic studies by the National Institute for Health and Care Excellence (10), which was adapted from an article by Hayden et al. (11). Included studies were awarded a score independently by 2 reviewers, and any discrepancies in the quality score were resolved by discussion, or by a third reviewer if no consensus could be reached.

Strategy for data synthesis

Results were collated thematically, with variables categorized as demographics, auxology, infections, physiological markers, and laboratory investigations. Quantitative data synthesis was undertaken if $\geq 20\%$ of studies provided data on the same factor from a multivariable model. We conducted a random-effects meta-analysis stratified by the measure of effect (HR, RR, or OR). No single measure of effect was given precedence; instead, inferences were drawn from all the available data. Tests were carried out using Stata statistical software version 16 (StataCorp LLC, 2019). We calculated a summary measure of effect, along with a 95% CI, and heterogeneity was assessed using Cochran's Q test and Higgins' I^2 test statistic.

Results

Of 1423 articles generated by the search, 28 fulfilled the inclusion criteria; 19 studies evaluated all children hospitalized with SAM, and 9 evaluated subgroups of children with SAM [diarrhea ($n = 4$), pneumonia ($n = 2$), kwashiorkor ($n = 1$), shigellosis ($n = 1$), and those with blood cultures ($n = 1$)]. **Supplemental Figure 1** shows the search process, including the number of included studies.

Full studies

Table 1 summarizes the 19 full studies including all children with SAM (12–30). All were from sub-Saharan Africa, across 8 different countries. The majority (15 of 19; 79%) were single-center studies. One pair of studies in Uganda (19, 20) and 1 pair in Ethiopia (26, 29) partially share a data set. The median number of children in each study was 400 (IQR: 120–545) and mean inpatient mortality was 15.7% (95% CI: 10.4%, 21.0%).

Supplemental Table 1 shows the risk of bias for each study. Ten studies (53%) were classed as having low risk of bias, 7 (37%) medium risk, and 2 (10%) high risk. The majority (11 of 19; 58%) of studies did not describe the causes of attrition from their cohort and/or failed to reduce selection bias in their studies.

TABLE 1 Risk factors independently associated with inpatient mortality in SAM¹

Study	Country	Year	Single/multi center	SAM diagnostic criteria	# of SAM Children	Age range	Inpatient mortality	HIV prevalence	Anthropometry			Infection			System Failures			Laboratory Investigations					Significant Factors associated with inpatient mortality on multivariable analysis					
									Sex: female	Age	Malnutrition anthropometry	Presence of edema	Hospital type	Family Demographics	HIV positive	Malaria	Clinical Sepsis/Shock	Blood Culture Positive	Lower GI problems	Respiratory Infection	Oral Thrush	Abnormal Obs or		Neurologically abnormal	Feeding/vomiting problems	Skin/eye problems	Electrolyte abnormalities	Blood counts abnormal
Babirekere-Iriso(12)	Uganda	2006	Single	WHO	134	6 - 59mo	20.1%	44.0%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Bacteremia ²
Bachou(13)	Uganda	2006	Single	WHO	220	0 - 59mo	24.0%	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Anemia needing transfusion ³	
Chimhuya(14)	Zimbabwe	2007	Single	WHO	244	0 - 59mo ⁴	41.4%	28.7%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Age < 18mo; W:H > 70%; Pneumonia; Diarrhea ³	
De Maayer(15)	South Africa	2011	Multi	WHO or Wellcome	113	0 - 59mo	11.5%	51.0%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	HIV; Pallor; Shock ²	
Irena(16)	Zambia	2011	Single	WHO	430	6 - 59mo	40.5%	38.6%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Diarrhea; Marasmic-kwashiorkor; Age 18-23mo ³	
Nhampossa(17)	Mozambique	2013	Single	WHO	1365	0 - 59mo	7.1%	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Breastfeeding ⁵ ; Prostration; Candidiasis; Hypoglycemia ⁶ ; Bacteremia ⁷ ; Malaria; Pallor ⁸ ; Edema ⁸ ; Diarrhea ⁸	
Vygent(18)	Niger	2013	Single	WHO	632	0 - 6mo	5.9%	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Infection ³	
Bartz(19)	Uganda	2014	Single	WHO	75	6 - 59mo	12.2%	24.0%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	HIV status; Leptin level ³	
Freemark(20)	Uganda	2015	Single	'severely malnourished'	75	6 - 59mo	12.5%	24.0%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Low leptin level ³	
Heilskov(21)	Uganda	2015	Single	WHO	120	6 - 59mo	N/A	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Lichenoid skin changes; Bullae/erosions ³	
Destaf(22)	Ethiopia	2015	Single	WHO	441	0 - 59mo	28.7%	4.1%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Sex ²	
Attia(23)	Malawi	2016	Single	WHO	79	6 - 60mo ³	17.7%	44.3%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Diarrhea; low SCFA; Cytokines ⁶	
Oumer(24)	Ethiopia	2016	Single	WHO	617	0 - 59mo	7.6%	2.9%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	HIV; Low appetite; Higher W:H Z; Malaria ²	
Ryter(25)	Uganda	2017	Single	WHO	120	6 - 59mo	14.0%	19.0%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Age >24mo; Higher MUAC; Severity of illness; Thrush; Impaired consciousness; Low phosphate ²	
Girum(26)	Ethiopia	2017	Multi	'national protocol' ⁷	545	0 - 59mo	9.3%	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Age <24mo; Altered pulse rate; Temperature; Shock; NG tube; Anemia; hypoglycemia; Facility: hospital ²	
Guesh(27)	Ethiopia	2018	Multi	WHO	569	0 - 59mo	3.7%	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Impaired consciousness; Urban residence ²	
Nabukeera-Baringir(28)	Uganda	2018	Single	WHO	400	6 - 59mo	9.8%	11.7%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Low oxygen saturations ²	
Girum(29)	Ethiopia	2018	Multi	'national protocol' ⁷	400	0 - 59mo	8.3%	--	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Altered pulse rate; needs NG; Anemia; Hypoglycemia ²	
Desyibelew(30)	Ethiopia	2019	Single	WHO	401	0 - 59mo	8.5%	6.5%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	Age < 24mo; Fully vaccinated; HIV; Heart failure ²	

¹ * < 24 mo only; ** >= 24 mo only; GI, gastrointestinal; MUAC, midupper arm circumference; N/A, not applicable; NG, nasogastric; Obs, observations; SAM, severe acute malnutrition; W:H, weight for height; †, increased independent risk with factor on multivariate analysis; (†), increased risk on univariate analysis but not multivariate analysis; ↔, analyzed, but not significant on multivariate analysis; ‡, decreased risk on univariate analysis but not multivariate analysis; †, increased independent risk with factor on multivariate analysis; ‡, decreased independent risk with factor on multivariate analysis.

² Calculated by Cox's multivariate analysis.

³ Calculated by multivariate logistic regression.

⁴ Confirmed with author.

⁵ The authors were contacted to clarify the upper age limit: the oldest children recruited were aged 59 completed months; the article was therefore eligible for inclusion.

⁶ Calculated by partial least squares modeling.

⁷ The Ethiopian national protocol is based on the WHO guidelines.

⁸ Calculated by Gompertz regression.

Sex, age, anthropometry, and edema.

Six studies evaluated child sex as a predictor of mortality. One study identified increased inpatient mortality among females [adjusted hazard ratio (aHR): 1.47; 95% CI: 1.01, 1.59] (22); 5 found no evidence of mortality difference by sex (13, 16, 25, 28, 30).

Five studies explored the impact of child age, using different age thresholds. There was generally an increased risk of mortality for children at either end of the 0- to 59-mo age range. One study in Zimbabwe found children aged <18 mo were at greater risk of inpatient mortality than those aged >18 mo [adjusted odds ratio (aOR): 2.27; 95% CI: 1.20, 4.29] (14), similarly to 2 studies in Ethiopia, which suggested those aged <24 mo had increased mortality [aHR: 2.84; 95% CI: 1.10, 7.34 (26), and aHR: 3.71; 95% CI: 1.37, 10.05 (30)]. A Zambian study found those aged 18–23 mo were at decreased risk of mortality (aOR: 0.44; 95% CI: 0.23, 0.85), whereas younger children were at increased risk [aOR: 1 (reference standard) (16)]. Rytter et al. (25) found that children aged 24–59 mo (aHR: 5.7; 95% CI: 1.8, 18.2) were at increased risk of dying compared with those aged 12–23 mo.

Anthropometry was assessed by MUAC and/or weight-for-height *z* score (WHZ). Children with a higher MUAC had lower mortality in 1 study (aHR: 0.6; 95% CI: 0.4, 1.0) (25), but no significant difference in outcomes in another study (aHR: 0.82; 95% CI: 0.64, 1.06) (28). Higher WHZ was associated with lower mortality in 2 studies [aHR: 0.95; 95% CI: 0.90, 0.99 (24), and aOR: 0.38; 95% CI: 0.18, 0.81 (14)]; 2 other studies showed no significant difference in mortality by WHZ (19, 26).

The effect of edema was heterogeneous across studies. Some found that the presence of edema increased the risk of mortality in children aged >24 mo (aOR: 10.77; 95% CI: 2.47, 46.7) (17), or in those with marasmic-kwashiorkor (aOR: 2.8; 95% CI: 1.52, 5.15) (16), but others indicated no evidence of difference by edema status (25, 28).

Demographics.

An urban compared with a rural setting was associated with higher inpatient mortality in studies from Ethiopia that investigated either urban or rural populations (aHR: 2.37; 95% CI: 1.12, 6.64) (27), or treatment in predominantly urban hospitals compared with more rural stabilization centers (aHR: 4.77; 95% CI: 1.64, 13.9) (29). Another Ethiopian study showed no evidence of increased mortality in urban children (aHR: 1.24; 95% CI: 0.51, 3.01) (24).

Co-infections.

The most commonly described co-infection was HIV, in 12 of 19 (63%) studies. Studies including HIV as a factor in multivariable analysis generally identified an increased risk of inpatient mortality among children with HIV. One study found a clear impact of HIV on mortality [aHR: 3.82; 95% CI: 1.30, 11.23 (30)]. Three studies reported a large effect size, but with imprecise estimates, likely due to smaller numbers of HIV-positive children [aHR: 6.1; 95% CI: 1.3, 28 (15); aOR: 117; 95% CI: 2, 6809 (19); aHR: 11.57; 95% CI: 2.34, 57.2 (24)]. Four studies had point estimates that suggested increased mortality in

children with HIV, but the 95% CIs included unity (13, 16, 25, 28).

There were heterogeneous findings for malaria, with 1 study from Mozambique showing that malaria was protective [aOR for <24 mo: 0.49; 95% CI: 0.26, 0.95; aOR for >24 mo: 0.3; 95% CI: 0.09, 0.92 (17)], and another from Ethiopia showing that it was deleterious [aHR: 12.7; 95% CI: 4.57, 35.3 (24)]. This difference may be explained by the very different prevalence of malaria in these studies: in Mozambique, 52% of the study population had malaria parasitemia (17), whereas in the Ethiopian study the figure was only 2.6% (24).

Children showing signs of serious illness due to infection generally had poorer outcomes. Those displaying clinical signs of shock had significantly higher mortality in 3 studies [aHR: 4.8; 95% CI: 1.5, 15 (15); aHR: 3.9; 95% CI: 1.4, 11.3 (25); aHR: 3.81; 95% CI: 1.83, 7.92 (26)] but there was no evidence of increased mortality in another [aHR: 2.0; 95% CI: 0.51, 7.84 (27)]. In addition, 1 study examining children with “infections” found no effect on mortality (18). Children with bacteremia had independently higher mortality than those without bacteremia [aOR: 8.45; 95% CI: 3.2, 22.3 (12); aOR for <24 mo: 1.95; 95% CI: 1.0, 3.78 (17)] in studies from Kenya and Mozambique. Children presenting with fever had a better outcome than those without fever in a Zambian study [aOR: 0.59; 95% CI: 0.37, 0.93 (16)]; there was no significant association between fever and mortality in 2 other studies [aHR: 0.3; 95% CI: 0.1, 1.2 (25); aHR: 1.6; 95% CI: 0.83, 3.07 (28)].

Diarrhea was independently associated with increased mortality in 3 studies [aOR: 3.42; 95% CI: 1.53, 7.65 (14); aOR: 2.5; 95% CI: 1.5, 4.1 (16); aOR for >24 mo: 4.39; 95% CI: 1.49, 12.9 (17)], although infectious etiology was not investigated. Three other studies showed no significant association between diarrhea and mortality (24, 25, 28). One study found that pneumonia was significantly associated with mortality [aOR: 2.21; 95% CI: 1.08, 4.52 (14)], whereas 3 others did not (24, 28, 30). Oral candidiasis was significantly associated with mortality in 2 studies [aHR: 5.0; 95% CI: 1.6, 15.2 (25), and aOR for >24 mo: 20.15; 95% CI: 2.97, 142 (17)]; HIV prevalence was only measured in 1 of these studies. In a Ugandan study, measles [aHR: 0.73; 95% CI: 0.27, 1.99 (30)] and tuberculosis [aHR: 0.91; 95% CI: 0.27, 3.11 (30)] were not independently associated with mortality. The same study found that being fully vaccinated was associated with significantly reduced mortality [aHR: 0.16; 95% CI: 0.07, 0.36 (30)].

End-organ dysfunction.

Factors indicating end-organ dysfunction were generally independently associated with increased inpatient mortality. An abnormal pulse rate was associated with increased mortality in 2 studies [aHR: 3.93; 95% CI: 1.58, 9.76 (26); aHR: 2.44; 95% CI: 1.47, 4.0 (29)], but an abnormal respiratory rate was not [aHR: 0.99; 95% CI: 0.93, 1.07 (28)]. Congestive heart failure was associated with elevated mortality in Ethiopia [aHR: 6.98; 95% CI: 2.42, 20.09 (30)]. Children with reduced consciousness on presentation consistently had higher mortality in 3 separate studies [aHR: 16.7; 95% CI: 3.1, 90.4 (25); aHR: 6.7; 95% CI: 2.43, 19.9 (27); aOR for <24 mo old: 3.2; 95% CI: 1.77, 5.8, and aOR for >24 mo old: 17.4; 95% CI: 4.4, 69.2 (17)].

Appetite.

Three studies found that lack of appetite was significantly associated with increased mortality [aHR: 2.75; 95% CI: 1.08, 6.99 (24); aHR: 3.18; 95% CI: 1.18, 8.58 (26); aHR: 1.8; 95% CI: 1.04, 3.1 (29)], whereas 2 studies showed no significant increase in mortality (25, 28).

Skin lesions.

Several studies evaluated skin lesions, which were defined differently between studies. Children with “lichenoid skin changes” had a significantly poorer outcome in 1 study from Uganda when adjusted for edema [measure of effect not given (21)], whereas in another Ugandan study “skin lesions” were significantly associated with mortality when adjusted for age and sex, but not in full multivariable analysis (28), including adjusting for edema status. An Ethiopian study found a nonsignificant trend for “skin ulcers” to be protective against mortality (27).

Laboratory results.

One study found no association between abnormal sodium or potassium and mortality, but did find that low phosphate on day 2 of admission (but not on day 1) was associated independently with increased mortality [aHR: 8.7; 95% CI: 2.5, 30.1 (25)]. Findings for anemia were heterogeneous. There was evidence that children with anemia, determined only by clinical criteria, had higher mortality in 3 studies [aHR: 1.53; 95% CI: 1.02, 2.3 (29); aHR: 5.1; 95% CI: 1.1, 24 (15); aOR for >24 mo old: 3.35; 95% CI: 1.05, 10.63 (17)], but there was no significant increase in mortality in another study (27). Where laboratory hemoglobin values were assessed as a continuous outcome, 2 studies found no significant association with mortality per unit change in hemoglobin (25, 28). Anemia requiring transfusion was significantly associated with mortality in 1 study [aOR: 5.1; 95% CI: 2.2, 12 (13)]. Abnormal white cell counts were not associated with mortality in 1 study (28).

Raised inflammatory markers were associated with mortality in several studies, which used different biomarkers and varying cutoffs. One Ugandan study found a raised C-reactive protein (CRP) concentration >15 mg/L was associated with increased mortality, compared with children with values ≤15 mg/L [aHR: 12.6; 95% CI: 1.6, 100 (25)]. Another study investigated CRP values >10 mg/L, but found no significant association with mortality [aHR: 4.0; 95% CI: 0.94, 17 (28)]. One study found that “systemic inflammation,” characterized by multiple cytokines including IL-6, TNF- α , and IL1 receptor antagonist, was independently associated with mortality, but did not report specific measures of effect for each cytokine (23). The same study found that intestinal inflammation, as measured by fecal calprotectin and fecal SCFAs, was associated with mortality (23). Higher plasma leptin was associated with reduced mortality in 1 study [aOR: 0.91; 95% CI: 0.83, 0.99 (19, 20)], but adiponectin was not [aOR: 1.0; 95% CI: 0.99, 1.0 (19)].

Substudies

Of the 9 studies evaluating subgroups of children admitted with SAM (31–39), two-thirds (6 of 9) were undertaken in

Bangladesh, and HIV prevalence was not reported. **Table 2** shows a summary of the variables significantly independently associated with mortality. Three studies at least partly shared a data set (36–38). Owing to the heterogeneity of the populations being selected using different characteristics, these substudies were not suitable for meta-analysis.

In the substudies, pneumonia or hypoxia was the commonest significant predictor of mortality, in a Malawian study of kwashiorkor [no effect size given, $P = 0.002$ (31)] and 3 studies from Bangladesh focused on shigellosis [aOR: 2.5; 95% CI: 1.1, 5.5 (33)], diarrhea [aOR: 3.0; 95% CI: 1.2, 7.3 (35)], and children admitted with pneumonia [aOR: 23.15; 95% CI: 4.38, 122.42 (36)]. Altered observations such as hypothermia were also significant predictors of mortality in 2 studies from Bangladesh [aOR: 5.7; 95% CI: 1.5, 22.1 (33) and aOR: 3.5; 95% CI: 1.3, 9.4 (35)].

Quantitative meta-analysis

Of the 57 variables measured across studies, 10 were included in ≥4 studies (20%) and were therefore considered for the quantitative data synthesis: HIV, anemia, diarrhea, edema, WHZ, shock, lack of appetite, pneumonia, age < 24 mo, and sex. Anemia was not included in the meta-analysis, because definitions across studies varied considerably: hemoglobin concentration <4 g/dL (13), <9.3 g/dL (14), or <11 g/dL (24); clinical pallor of mucous membranes (15, 17, 26, 27, 29); or a value of difference in risk per unit change in hemoglobin (25, 28). Nine variables were therefore included in the meta-analysis.

Figure 1 shows the results of the meta-analysis for sex, edema, WHZ, HIV infection, and age < 24 mo. All factors showed some evidence of heterogeneity. Of the 6 studies including child sex, only 1 found a significant association with mortality (22). In meta-analysis, the HR (0.87; 95% CI: 0.42, 1.79) and OR (0.98; 95% CI: 0.65, 1.47) demonstrated no evidence of association between sex and mortality (**Figure 1A**). There was strong evidence of an association between presence of edema and higher mortality in the 3 studies reporting ORs (OR: 2.43; 95% CI: 1.24, 4.77), but not in the 2 studies reporting HRs (HR: 0.85; 95% CI: 0.49, 1.48) (**Figure 1B**). Overall, higher WHZs were associated with lower mortality in the 2 studies reporting ORs (OR: 0.44; 95% CI: 0.24, 0.80), but not in the 2 studies reporting HRs (HR: 0.87; 95% CI: 0.66, 1.14) (**Figure 1C**). Eight studies including HIV as an explanatory variable were included in the meta-analysis. The 5 studies reporting HRs found >4-fold higher mortality among HIV-positive than among HIV-negative children (HR: 4.32; 95% CI: 2.31, 8.08); by contrast, the 3 studies reporting ORs found no significant increase in mortality in HIV-positive children (OR: 2.03; 95% CI: 0.88, 4.67) (**Figure 1D**). In the 4 studies specifically exploring an age cutoff of 24 mo, there was no significant association between age and mortality in the meta-analysis (**Figure 1E**).

Figure 2 shows the results of the meta-analysis for factors assessing acute clinical illness during hospitalization. The presence of diarrhea was associated with almost 3-fold higher inpatient mortality (HR: 2.84; 95% CI: 1.40, 5.75; OR: 2.91; 95% CI: 1.96, 4.33) (**Figure 2A**). Shock, as judged by clinical criteria, was associated with almost 4-fold higher mortality (HR: 3.67; 95% CI: 2.24, 6.03) (**Figure 2B**). Pneumonia was associated with 2-fold higher mortality in the meta-analysis (HR: 1.89; 95% CI:




FIGURE 1 Forest plots of the meta-analysis of sex (A), presence of edema (B), WHZ (C), HIV infection (D), and age <2 y (E), on mortality in children admitted with SAM. Results with HRs are shown in red and studies reporting results in ORs are shown in green. ^aThis result was reported as aHR: 0.679 (95% CI: 0.63, 0.99) among males. However, given the CI presented, it is likely there is a typographical error and the point estimate should be 0.79. We contacted the authors but did not receive a reply. We have therefore used the value as published. We found no change in inference from using the likely alternative value. WHZ, weight-for-height z score.

1.19, 3.02; OR: 2.21; 95% CI: 1.08, 4.52) (Figure 2C). Lacking appetite and/or requiring a nasogastric (NG) tube for feeding was associated with increased mortality in a meta-analysis of 5 studies (HR: 2.16; 95% CI: 1.48, 3.16) (Figure 2D).

Discussion

In this first (to our knowledge) published meta-analysis of the factors independently associated with mortality among children hospitalized with SAM, we aimed to identify those children who were most vulnerable at admission for inpatient management. The mean inpatient mortality in this review was 15.7%, which is well above global targets of 5%–10% (40, 41), emphasizing the pressing need to reduce hospital deaths. We identified 6 statistically independent factors associated with increased mortality: low WHZ, HIV infection, diarrhea, pneumonia, shock, and lack of appetite/requiring NG feeding. These factors reflect the severity and type of SAM, together with comorbidities and decompensation of physiological pathways underlying this complex multisystem condition,

thereby highlighting potential points for intervention to improve outcomes.

SAM presents as 2 distinct clinical phenotypes: nonedematous (previously called marasmus) and edematous (previously called kwashiorkor) (42). Nomenclature and classifications have changed over time, with the 1970 Wellcome classification (based on weight-for-age) further distinguishing the clinical concepts of marasmus, marasmic-kwashiorkor, and kwashiorkor (43). Waterlow (44) in 1972 distinguished the acute process of wasting in a classification based on weight-for-height. In our systematic review, children with lower WHZs at admission were at higher risk of mortality in most studies, and in meta-analysis higher WHZs were associated with lower mortality (OR: 0.44; 95% CI: 0.24, 0.80). Lower WHZ reflects a greater degree of wasting, which is characterized by a loss of muscle mass and subcutaneous fat (45), together with underlying physiological derangement. Edematous SAM is characterized by bilateral pitting edema, enlarged steatotic liver, skin and hair changes, and apathy (46). Studies report heterogeneous results for the effects of edema on mortality. In a meta-analysis stratified by the measure of




FIGURE 2 Forest plots of the meta-analysis of diarrhea (A), presence of shock (B), pneumonia (C), and lack of appetite/needling NG feeding (D), on mortality in children admitted with SAM. Results with HRs are shown in red and studies reporting results in ORs are shown in green. NG, nasogastric.

effect in each study, the presence of edema was associated with higher odds of death (OR: 2.43; 95% CI: 1.24, 4.77), but not with a higher hazard of death (HR: 0.85; 95% CI: 0.49, 1.48). It is therefore difficult to conclude what impact edema has on mortality among children with complicated SAM. Over 70 y after it was first described (42), the mechanisms underlying edema in malnutrition remain poorly understood. The massive increase in liver fat observed suggests that distinct metabolic processes occur in edematous compared with nonedematous SAM. There is increasing interest in whether perturbations in the gut microbiome underlie fatty liver changes and generation of edema (47). Those with edema also have differences in DNA methylation compared with those without edema, but the significance of these epigenetic changes or the patterns of genes involved is unclear (48). The mixed clinical presentation of marasmic-kwashiorkor was shown by Irena et al. (16) to have a higher mortality than kwashiorkor alone, suggesting that the combination of low WHZ and edema may increase mortality risk, but further studies from other settings are needed to clarify the outcomes of children with and without edema.

HIV infection emerged as a clear risk factor for mortality in meta-analysis, with >4-fold higher mortality than among HIV-negative children (HR: 4.32; 95% CI: 2.31, 8.08), consistent with a previous systematic review (49). Although prevention of mother-to-child transmission interventions have reduced the proportion of children acquiring HIV, the mean HIV prevalence in this systematic review was 24.9%. Many of the included studies were undertaken before widespread availability of antiretroviral therapy (ART), although recent data from Zimbabwe and Zambia highlight 4-fold higher mortality among HIV-positive children with SAM, despite the availability of ART (50). HIV has influenced the epidemiology, presentation, and outcomes of children with SAM (51). A consultation in 2004 recognized the need to better understand the impact of implementing WHO guidelines for nutritional rehabilitation on the case fatality rate

among HIV-positive children (52). Children with HIV and SAM are less likely to have edema, and more likely to have presentations of advanced HIV disease, such as fungal infections and persistent diarrhea (53). HIV infection likely compounds SAM by exacerbating immunosuppression, inflammation, and enteropathy (54), although there is a recognized need to better understand the underlying pathophysiology (51) and to identify management approaches for children with coexistent HIV and SAM.

Shock at admission, representing a state of decompensation and end-organ dysfunction, was associated with an almost 4-fold increased risk of inpatient death (HR: 3.67; 95% CI: 2.24, 6.03). Two studies not eligible for meta-analysis (26, 29) similarly found that an abnormal pulse rate was associated with death. Once children with SAM have progressed to shock, their poor physiological reserve and high risk of fluid overload during resuscitation likely contribute to the poor prognosis. Diarrhea at admission was associated with a 3-fold increased risk of inpatient death (OR: 2.91; 95% CI: 1.96, 4.33). Diarrhea is a prominent feature of SAM, especially as the severity of malnutrition increases, due to a vicious cycle of underlying pathology, particularly in children with HIV (55). Given the importance of clinical and subclinical enteropathy in driving mortality, recent (56) and ongoing studies (57) are exploring the role of additional therapeutic interventions to heal the gut during inpatient management of complicated SAM. Pneumonia complicating SAM was associated with a 2-fold increased risk of death (HR: 1.89; 95% CI: 1.19, 3.02). Pneumonia in children with reduced physiological reserve results in even greater metabolic demands to overcome fever, increased cardiac output, work of breathing, and hypoxemia. In children with SAM and a diagnosis of pneumonia both the presence of severe sepsis (58) and hypoxemia (59) have been shown to greatly increase the risk of death. Children with a loss of appetite or requiring NG feeds had a 2-fold increased risk of death (HR: 2.16; 95% CI:

1.48, 3.16). This finding could reflect the extent of the underlying pathology of SAM, which results in apathy and loss of appetite, or the severity of the child's clinical condition, because those with critical illness may have an altered level of consciousness and an unsafe swallow. Although NG feeding is an essential intervention during therapeutic rehabilitation, the need for an NG tube should also be recognized by clinicians as a surrogate marker of mortality risk. Three studies not eligible for meta-analysis showed that reduced consciousness was associated with higher mortality. A recent meta-analysis of inpatient treatment outcomes among children with SAM in Ethiopia found diarrhea, dehydration, and anemia to be statistically significant predictors of inpatient mortality (60). This previous meta-analysis included only Ethiopian studies, did not restrict analyses to independent predictors of mortality, made no assessment of HIV, and 5 out of 21 of its open-access articles were not indexed in PubMed/Embase. Nevertheless, both the previous and the current meta-analysis suggest that markers of comorbidity (such as diarrhea) and physiological dysfunction are associated with elevated mortality.

Age and sex were not significant predictors of mortality in our meta-analysis. Although not eligible for meta-analysis, lower anthropometry measures and anemia also appeared to be associated with increased mortality in our narrative synthesis. MUAC, recognized as a validated measure in the 2013 update of the WHO guidelines, has relatively recently been introduced and was frequently not reported in earlier studies. Owing to its collinearity with WHZ, it may not be identified as a statistically independent determinant of mortality in multivariable analyses which include both variables (26), although it should be noted that some studies did not identify any association between MUAC and death even when WHZ was not included (23, 24, 28). Anemia was excluded from meta-analysis because there were considerable variations in the definitions used between studies. WHO estimates that 42% of children <5 y of age globally are anemic (61). Children in this age group with SAM are at particular risk of anemia, owing to increased iron demands during rapid growth; iron deficiency arising from inadequate diets; infections such as malaria (62), HIV, and hookworm (63); and micronutrient deficiencies. Iron deficiency negatively affects innate and adaptive immunity, further increasing vulnerability to infections (64–66). Our narrative review indicated heterogeneity between studies reporting associations between anemia (or hemoglobin concentrations) and mortality; further studies may help to clarify these relations.

Factors associated with mortality in >1 study, but ineligible for meta-analysis, included raised inflammatory markers, reduced consciousness, urban setting, absence of fever, presence of skin changes, and bacteremia. It is difficult to interpret the findings regarding study setting, because there was no apparent correction for the plausible selection bias among children admitted to an urban referral hospital, who are likely to be sicker. Fever, although a widely measured variable, may not be a good predictor of mortality because children with SAM frequently do not manifest the usual clinical signs of infection such as fever, potentially owing to impairment of the acute inflammatory response (67).

Our review has strengths and weaknesses. We set out to identify all relevant English-language studies since the 1999 WHO guidelines for management of SAM were published. The

studies included in our review were almost exclusively from sub-Saharan Africa; the only studies from South Asia, which has the largest number of children with SAM, focused on specific subgroups of children. We identified >50 baseline factors across studies, but definitions often varied, and only 9 variables were eligible for meta-analysis. For many of these variables, there was moderate to high heterogeneity, and it was difficult to ascertain the impact of edema in particular on mortality. The risk of bias among studies was assessed by evaluating domains of participation, attrition, prognostic factors, outcomes, confounding, and analysis. Of the 19 studies, 2 had a high risk of bias and 7 had a medium risk of bias. One study with a high risk of bias was included in the meta-analysis (27) of shock. This retrospective study had a high attrition rate (15.7% lost to follow-up) and unclear methods for measurement of each variable, imputation of missing data, and adjustment for confounding. The majority of studies with medium-to-high risk of bias failed to adequately describe the study sample, account for attrition, or describe those lost to follow-up. Several studies had limitations, including omitting results for factors that were not significant, which meant they could not be included in our meta-analysis; although 1 author responded to requests for additional data, we were unable to access the majority of data. In addition, studies reported different measures of effect, making a single meta-analysis for each factor difficult. We chose to stratify the results by measure of effect, as in previous meta-analyses (68), but this potentially limited our inferences, because not all data were combined into a single pooled effect. Finally, we chose to include only results from multivariable analysis, to identify statistically independent predictors of mortality, but it is likely that there are pathophysiological interdependencies between variables, meaning we may have missed some clinically valuable (but not statistically independent) factors associated with mortality—a limitation noted by some other authors (69).

In summary, our meta-analysis identified 6 independent factors associated with mortality in children hospitalized for complicated SAM: WHZ, HIV infection, diarrhea, pneumonia, shock, and lack of appetite/requiring NG feeding. Despite adherence to WHO guidelines it has been demonstrated that mortality remains unacceptably high in children hospitalized with SAM (69). Early recognition of these prognostic factors in the community, with referral to inpatient facilities, and risk stratification at hospital admission may help to reduce inpatient mortality among children with SAM. The independent factors we identified reflect the interaction of predisposing factors, comorbidities, and end-organ manifestations of SAM, which ultimately drive mortality in these critically ill children. We require a greater mechanistic understanding of the underlying causal pathways, to ascertain whether early interventions targeting these factors may reduce mortality in this high-risk population.

The authors' responsibilities were as follows—RK and AJP: conceived the project; RK, JPS, and RP: performed the searches and data extraction; JPS and RK: conducted the analysis; AJP: had oversight of the project; and all authors: drafted the paper and read and approved the final manuscript. The authors report no conflicts of interest.

References

1. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, Ezzati M, Grantham-McGregor S, Katz J, Martorell R, et al. Maternal

- and child undernutrition and overweight in low-income and middle-income countries. *Lancet* 2013;382(9890):427–51.
2. Rytter MJ, Kolte L, Briend A, Friis H, Christensen VB. The immune system in children with malnutrition—a systematic review. *PLoS One* 2014;9(8):e105017.
 3. Bourke CD, Berkley JA, Prendergast AJ. Immune dysfunction as a cause and consequence of malnutrition. *Trends Immunol* 2016;37(6):386–98.
 4. World Health Organization. The treatment and management of severe protein-energy malnutrition. Geneva, Switzerland: WHO; 1981.
 5. World Health Organization. Management of severe malnutrition: a manual for physicians and other senior health workers. Geneva, Switzerland: WHO; 1999.
 6. World Health Organization. Guidelines for the inpatient treatment of severely malnourished children. New Delhi, India: WHO Regional Office for South-East Asia; 2003.
 7. World Health Organization. Guideline: updates on the management of severe acute malnutrition in infants and children. Geneva, Switzerland: WHO; 2013.
 8. Williams P, Berkley J. Severe acute malnutrition update: current WHO guidelines and the WHO essential medicine list for children. *J Clin Pharmacol* 2016;56(11):1344–54.
 9. Tickell KD, Denno DM. Inpatient management of children with severe acute malnutrition: a review of WHO guidelines. *Bull World Health Organ* 2016;94(9):642–51.
 10. National Institute for Health and Care Excellence (NICE). Appendix I: methodology checklist: prognostic studies [Internet]. London: NICE; 2012. Available from: <https://www.nice.org.uk/process/pmg6/resources/the-guidelines-manual-appendices-bi-2549703709/chapter/appendix-i-methodology-checklist-prognostic-studies>, [accessed 31 March, 2020].
 11. Hayden JA, Cote P, Bombardier C. Evaluation of the quality of prognosis studies in systematic reviews. *Ann Intern Med* 2006;144(6):427–37.
 12. Babirekere-Iriso E, Musoke P, Kekitiinwa A. Bacteraemia in severely malnourished children in an HIV-endemic setting. *Ann Trop Paediatr* 2006;26(4):319–28.
 13. Bachou H, Tumwine JK, Mwadime RKN, Tylleskär T. Risk factors in hospital deaths in severely malnourished children in Kampala, Uganda. *BMC Pediatr* 2006;6:7.
 14. Chimhuya S, Kambarami RA, Mujuru H. The levels of malnutrition and risk factors for mortality at Harare Central Hospital-Zimbabwe: an observation study. *Cent Afr J Med* 2007;53(5–8):30–4.
 15. De Maayer T, Saloojee H. Clinical outcomes of severe malnutrition in a high tuberculosis and HIV setting. *Arch Dis Child* 2011;96(6):560–4.
 16. Irena AH, Mwambazi M, Mulenga V. Diarrhea is a major killer of children with severe acute malnutrition admitted to inpatient set-up in Lusaka, Zambia. *Nutr J* 2011;10:110.
 17. Nhampossa T, Sigauque B, Machevo S, Macete E, Alonso P, Bassat Q, Menendez C, Fumado V. Severe malnutrition among children under the age of 5 years admitted to a rural district hospital in southern Mozambique. *Public Health Nutr* 2013;16(9):1565–74.
 18. Vygen SB, Roberfröid D, Captier V, Kolsteren P. Treatment of severe acute malnutrition in infants aged <6 months in Niger. *J Pediatr* 2013;162(3):e15–21.e3.
 19. Bartz S, Mody A, Hornik C, Bain J, Muehlbauer M, Kiyimba T, Kiboneka E, Stevens R, Bartlett J, St Peter JV, et al. Severe acute malnutrition in childhood: hormonal and metabolic status at presentation, response to treatment, and predictors of mortality. *J Clin Endocrinol Metab* 2014;99(6):2128–37.
 20. Freemark M. Metabolomics in nutrition research: biomarkers predicting mortality in children with severe acute malnutrition. *Food Nutr Bull* 2015;36(1 Suppl):S88–92.
 21. Heilskov S, Vestergaard C, Babirekere E, Ritz C, Namusoke H, Rytter M, Deleuran M. Characterization and scoring of skin changes in severe acute malnutrition in children between 6 months and 5 years of age. *J Eur Acad Dermatol Venereol* 2015;29(12):2463–9.
 22. Desta KS. Survival status and predictors of mortality among children aged 0–59 months with severe acute malnutrition admitted to Stabilization Center at Sekota Hospital Waghemra Zone. *J Nutr Disorders Ther* 2015;5:160.
 23. Attia S, Versloot CJ, Voskuil W, van Vliet SJ, Di Giovanni V, Zhang L, Richardson S, Bourdon C, Netea MG, Berkley JA, et al. Mortality in children with complicated severe acute malnutrition is related to intestinal and systemic inflammation: an observational cohort study. *Am J Clin Nutr* 2016;104(5):1441–9.
 24. Oumer A, Mesfin F, Demena M. Survival status and predictors of mortality among children aged 0–59 months admitted with severe acute malnutrition in Dilchora Referral Hospital, Eastern Ethiopia, EAJHB. 2016;1(1):13–22.
 25. Rytter MJ, Babirekere-Iriso E, Namusoke H, Christensen VB, Michaelsen KF, Ritz C, Mortensen CG, Mupere E, Friis H. Risk factors for death in children during inpatient treatment of severe acute malnutrition: a prospective cohort study. *Am J Clin Nutr* 2017;105(2):494–502.
 26. Girum T, Kote M, Tariku B, Bekele H. Survival status and predictors of mortality among severely acute malnourished children <5 years of age admitted to stabilization centers in Gedeo Zone: a retrospective cohort study. *Ther Clin Risk Manag* 2017;13:101–10.
 27. Guesh G, Degu G, Abay M, Beyene B, Brhane E, Brhane K. Survival status and predictors of mortality among children with severe acute malnutrition admitted to general hospitals of Tigray, North Ethiopia: a retrospective cohort study. *BMC Res Notes* 2018;11(1):832.
 28. Nabukeera-Barungi N, Grenov B, Lanyero B, Namusoke H, Mupere E, Christensen VB, Michaelsen KF, Mølgaard C, Rytter MJ, Friis H. Predictors of mortality among hospitalized children with severe acute malnutrition: a prospective study from Uganda. *Pediatr Res* 2018;84(1):92–8.
 29. Girum T, Muktar E, Worku A. Comparative analysis of the survival status and treatment outcome of under-five children admitted with severe acute malnutrition among hospital-based and health center based stabilization centers, south Ethiopia. *Open Public Health J* 2018;11(1):209.
 30. Desyibelew HD, Baraki AG, Dadi AF. Mortality rate and predictors of time to death in children with severe acute malnutrition treated in Felege-Hiwot Referral Hospital Bahir Dar, Northwest Ethiopia. *BMC Res Notes* 2019;12(1):409.
 31. Manary MJ, Brewster DR. Intensive nursing care of kwashiorkor in Malawi. *Acta Paediatr* 2000;89(2):203–7.
 32. Amadi B, Kelly P, Mwiya M, Mulwazi E, Sianongo S, Changwe F, Thomson M, Hachungula J, Watuka A, Walker-Smith J, et al. Intestinal and systemic infection, HIV, and mortality in Zambian children with persistent diarrhea and malnutrition. *J Pediatr Gastroenterol Nutr* 2001;32(5):550–4.
 33. van den Broek JM, Roy SK, Khan WA, Ara G, Chakraborty B, Islam S, Banu B. Risk factors for mortality due to shigellosis: a case-control study among severely-malnourished children in Bangladesh. *J Health Popul Nutr* 2005;23(3):259–65.
 34. Chisti MJ, Ahmed T, Bardhan PK, Salam MA. Evaluation of simple laboratory investigations to predict fatal outcome in infants with severe malnutrition presenting in an urban diarrhoea treatment centre in Bangladesh. *Trop Med Int Health* 2010;15(11):1322–5.
 35. Roy SK, Buis M, Weersma R, Khatun W, Chowdhury S, Begum A, Sarker D, Thakur SK, Khanam M. Risk factors of mortality in severely-malnourished children hospitalized with diarrhoea. *J Health Popul Nutr* 2011;29(3):229–35.
 36. Chisti MJ, Salam MA, Ashraf H, Faruque ASG, Bardhan PK, Hossain MI, Shahid ASMSB, Shahunja KM, Das SK, Imran G, et al. Clinical risk factors of death from pneumonia in children with severe acute malnutrition in an urban critical care ward of Bangladesh. *PLoS One* 2013;8(9):e73728.
 37. Chisti MJ, Salam MA, Bardhan PK, Faruque ASG, Shahid ASMSB, Shahunja KM, Das SK, Hossain MI, Ahmed T. Treatment failure and mortality amongst children with severe acute malnutrition presenting with cough or respiratory difficulty and radiological pneumonia. *PLoS One* 2015;10(10):e0140327.
 38. Chisti MJ, Shahid AS, Shahunja KM, Bardhan PK, Faruque ASG, Shahrin L, Das SK, Barua DK, Hossain MI, Ahmed T. Ileus in children presenting with diarrhea and severe acute malnutrition: a chart review. *PLoS Negl Trop Dis* 2017;11(5):e0005603.
 39. Obiero CW, Seale AC, Jones K, Ngari M, Bendon CL, Morpeth S, Mohammed S, Mturi N, Fegan G, Berkley JA. Should first-line empiric treatment strategies cover coagulase-negative staphylococcal infections in severely malnourished or HIV-infected children in Kenya? *PLoS One* 2017;12(8):e0182354.
 40. Ashworth A, Khanum S, Jackson A, Schofield C. Guidelines for the inpatient treatment of severely malnourished children. Geneva, Switzerland: World Health Organization; 2003.

41. Sphere Association. The Sphere handbook: humanitarian charter and minimum standards in humanitarian response. 4th ed. Geneva, Switzerland: Sphere Association; 2018.
42. Williams CD. Kwashiorkor. *J Am Med Assoc* 1953;153(14):1280–5.
43. Wellcome Trust Working Party. Classification of infantile malnutrition. *Lancet* 1970;2:302–3.
44. Waterlow J. Classification and definition of protein-calorie malnutrition. *BMJ* 1972;3(5826):566.
45. Williams CD. A nutritional disease of childhood associated with a maize diet. *Arch Dis Child* 1933;8(48):423–33.
46. Williams CD, Oxon BM, Lond H. Kwashiorkor: a nutritional disease of children associated with a maize diet. 1935. *Bull World Health Organ* 2003;81(12):912–13.
47. Kristensen KH, Wiese M, Rytter MJ, Ozcam M, Hansen LH, Namusoke H, Friis H, Nielsen DS. Gut microbiota in children hospitalized with oedematous and non-oedematous severe acute malnutrition in Uganda. *PLoS Negl Trop Dis* 2016;10(1):e0004369.
48. Schulze KV, Swaminathan S, Howell S, Jajoo A, Lie NC, Brown O, Sadat R, Hall N, Zhao L, Marshall K, et al. Edematous severe acute malnutrition is characterized by hypomethylation of DNA. *Nat Commun* 2019;10(1):5791.
49. Fergusson P, Tomkins A. HIV prevalence and mortality among children undergoing treatment for severe acute malnutrition in sub-Saharan Africa: a systematic review and meta-analysis. *Trans R Soc Trop Med Hyg* 2009;103(6):541–8.
50. Bwakura-Dangarembizi M, Amadi B, Dumbura C, Majo F, Ngosa D, Chasekwa B, Nathoo K, Walker S, Kelly MP, Prendergast AJ. One-year mortality in HIV-positive and HIV-negative children with complicated severe acute malnutrition. 16th Scientific Meeting of the Commonwealth Association of Paediatric Gastroenterology & Nutrition; 1–3 October 2019; Toronto, Canada.
51. Heikens GT, Bunn J, Amadi B, Manary M, Chhagan M, Berkley JA, Rollins N, Kelly P, Adamczick C, Maitland K, et al. Case management of HIV-infected severely malnourished children: challenges in the area of highest prevalence. *Lancet* 2008;371(9620):1305–7.
52. World Health Organization. Severe malnutrition: report of a consultation to review current literature, 6–7 September 2004. Geneva, Switzerland: WHO; 2005.
53. Bunn J, Thindwa M, Kerac M. Features associated with underlying HIV infection in severe acute childhood malnutrition: a cross sectional study. *Mal Med J* 2009;21(3):108–12.
54. Prendergast A, Kelly P. Enteropathies in the developing world: neglected effects on global health. *Am J Trop Med Hyg* 2012;86(5):756–63.
55. Prendergast AJ, Kelly P. Interactions between intestinal pathogens, enteropathy and malnutrition in developing countries. *Curr Opin Infect Dis* 2016;29(3):229–36.
56. Jones KDJ, Hünten-Kirsch B, Laving AMR, Munyi CW, Ngari M, Mikusa J, Mulongo MM, Odera D, Nassir HS, Timbwa M, et al. Mesalazine in the initial management of severely acutely malnourished children with environmental enteric dysfunction: a pilot randomized controlled trial. *BMC Med* 2014;12(1):133.
57. Kelly P, Bell L, Amadi B, Bwakura-Dangarembizi M, VanBuskirk K, Chandwe K, Chipunza M, Ngosa D, Chulu N, Hill S, et al. TAME trial: a multi-arm phase II randomised trial of four novel interventions for malnutrition enteropathy in Zambia and Zimbabwe - a study protocol. *BMJ Open* 2019;9(11):e027548.
58. Chisti MJ, Salam MA, Bardhan PK, Faruque ASG, Shahid ASMSB, Shahunja KM, Das SK, Hossain MI, Ahmed T, Borrow R. Severe sepsis in severely malnourished young Bangladeshi children with pneumonia: a retrospective case control study. *PLoS One* 2015;10(10):e0139966.
59. Chisti MJ, Salam MA, Ashraf H, Faruque ASG, Bardhan PK, Shahid ASMSB, Shahunja KM, Das SK, Ahmed T. Predictors and outcome of hypoxemia in severely malnourished children under five with pneumonia: a case control design. *PLoS One* 2013;8(1):e51376.
60. Wagnev F, Dessie G, Takele WW, Tadesse A, Islam SMS, Mulugeta H, Haile D, Negesse A, Abajobir AA. A meta-analysis of inpatient treatment outcomes of severe acute malnutrition and predictors of mortality among under-five children in Ethiopia. *BMC Public Health* 2019;19(1):1175.
61. World Health Organization. Prevalence of anaemia in children under 5 years (%), 2017 [Internet]. Geneva, Switzerland: WHO; [cited May 2020]. Available from: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-anaemia-in-children-under-5-years\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-anaemia-in-children-under-5-years(-)).
62. Menendez C, Fleming AF, Alonso PL. Malaria-related anaemia. *Parasitol Today* 2000;16(11):469–76.
63. Jonker FA, Calis JC, Phiri K, Brienen EA, Khoffi H, Brabin BJ, Verweij JJ, van Hensbroek MB, van Lieshout L. Real-time PCR demonstrates *Ancylostoma duodenale* is a key factor in the etiology of severe anemia and iron deficiency in Malawian pre-school children. *PLoS Negl Trop Dis* 2012;6(3):e1555.
64. Kumar V, Choudhry VP. Iron deficiency and infection. *Indian J Pediatr* 2010;77(7):789–93.
65. Drakesmith H, Prentice AM. Hcpidin and the iron-infection axis. *Science* 2012;338(6108):768–72.
66. Galan P, Thibault H, Preziosi P, Hercberg S. Interleukin 2 production in iron-deficient children. *Biol Trace Elem Res* 1992;32:421–6.
67. Christie C, Heikens GT, McFarlane D. Nosocomial and community-acquired infections in malnourished children. *J Trop Med Hyg* 1988;91(4):173–80.
68. Ardura-Garcia C, Stolbrink M, Zaidi S, Cooper PJ, Blakey JD. Predictors of repeated acute hospital attendance for asthma in children: a systematic review and meta-analysis. *Pediatr Pulmonol* 2018;53(9):1179–92.
69. Maitland K, Berkley JA, Shebbe M, Peshu N, English M, Newton CRJC. Children with severe malnutrition: can those at highest risk of death be identified with the WHO protocol? *PLoS Med* 2006;3(12):e500.